

Drawing 3D: Coordinates

**CSCI 4229/5229
Computer Graphics
Fall 2009**

The Issue

- Lorenz attractor (x,y,z) values are generally in the range (-50,50) in all three dimensions
- How do you get OpenGL to display the whole range of values?
- Assume
 - Lorenz values are in an array $x[i], y[i][z[i]], i=0, \dots, n-1$
 - double *dim* is defined somewhere as 50

Option 1

```
void display()
{
    for (i=0;i<n;i++)
        glVertex3d(x[i]/dim , y[i]/dim , z[i]/dim);
}
```

Option 2

```
void display()
{
    for (i=0;i<n;i++)
        glVertex4d(x[i] , y[i] , z[i] , dim);
}
```

Option 3

```
void display()
{
    glScaled(1/dim , 1/dim , 1/dim);
    for (i=0;i<n;i++)
        glVertex3d(x[i] , y[i] , z[i]);
}
```

Option 4

```
void display()
{
    for (i=0;i<n;i++)
        glVertex3d(x[i] , y[i] , z[i]);
}

void reshape()
{
    glScaled(1/dim , 1/dim , 1/dim);
}
```

Option 5

```
void display()
```

```
{...
```

```
    for (i=0;i<n;i++)
```

```
        glVertex3d(x[i] , y[i] , z[i]);
```

```
...}
```

```
void reshape()
```

```
{...
```

```
    glOrtho(-asp*dim,asp*dim,-dim,dim,-dim,dim);
```

```
...}
```